

**Note to Readers:** If you need assistance accessing items in this Supplemental Material, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Table of Contents for Supplemental Material**

### **Association of Prenatal Exposure to Persistent Organic Pollutants with Obesity and Cardiometabolic Traits in Early Childhood: The Rhea Mother–Child Cohort (Crete, Greece)**

Marina Vafeiadi, Vaggelis Georgiou, Georgia Chalkiadaki, Panu Rantakokko, Hannu Kiviranta,  
Marianna Karachaliou, Eleni Fthenou, Maria Venihaki, Katerina Sarri, Maria Vassilaki, Soterios

A. Kyrtopoulos, Emily Oken, Manolis Kogevinas, and Leda Chatzi

**Table S1.** Maternal and child characteristics of participants and non-participants.

**Table S2.** Maternal and child characteristics by concentrations of first trimester maternal serum POPs levels (pg/ml), Mother-child cohort “Rhea” in Crete, Greece (n=689).

**Table S3.** Associations of in utero POPs exposure with child lipids and adiponectin levels at 4 years of age, Mother-child cohort “Rhea” in Crete, Greece.

**Table S4.** Association between POPs and birth outcomes in multipollutant models including DDE, HCB and PCBs.

**Table S5.** Associations of in utero POPs exposure with child BMI at 4 years of age in the subgroups defined by infant sex, maternal smoking and maternal prepregnancy BMI status.

**Table S6.** Associations of in utero POP exposures with the outcomes under study in the total population and without out preterm births and low birth weight neonates.